

Managing Dental Erosion: Current Understanding and Future Directions

Course Author(s): Donald J. White, PhD; Warden H. Noble, DDS, MS, MEd; Nicola West, BDS, FDSRCS, PhD FDS (Rest Dent)

CE Credits: 2 hours

Intended Audience: Dentists, Dental Hygienists, Dental Assistants, Dental Students, Dental Hygiene Students, Dental Assistant Students

Date Course Online: 04/04/2017

Last Revision Date: N/A

Course Expiration Date: 04/03/2020

Cost: Free

Method: Self-instructional

AGD Subject Code(s): 10, 430, 730

Online Course: www.dentalcare.com/en-us/professional-education/ce-courses/ce517



Disclaimer: Participants must always be aware of the hazards of using limited knowledge in integrating new techniques or procedures into their practice. Only sound evidence-based dentistry should be used in patient therapy.

Introduction

The purpose of this course is to provide information on the dental erosion process, highlight key similarities and differences between dental erosion and caries, provide effective strategies to help educate patients who are at risk for developing dental erosion, and manage patients who are already experiencing the condition to minimize its effects.


Acknowledgements

To Mr. Robert V. Faller for assistance developing the course.

ADA CERP

The Procter & Gamble Company is an ADA CERP Recognized Provider.

ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of continuing dental education. ADA CERP does not approve or endorse individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry.

Concerns or complaints about a CE provider may be directed to the  | Continuing Education Recognition Program provider or to ADA CERP at: <http://www.ada.org/cerp>

Approved PACE Program Provider



The Procter & Gamble Company is designated as an Approved PACE Program Provider by the Academy of General Dentistry. The formal continuing education programs of this program provider are accepted by AGD for Fellowship, Mastership, and Membership Maintenance Credit. Approval does not imply acceptance by a state or provincial board of dentistry or AGD endorsement. The current term of approval extends from 8/1/2013 to 7/31/2017.

Provider ID# 211886

Conflict of Interest Disclosure Statement

- Dr. White is an employee of Procter & Gamble.
- Dr. Noble and Dr. West have done consulting work and given scientific lectures on behalf of P&G.

Course Contents

- Overview
- Learning Objectives
- Background
- Etiological Factors
 - Differences Between Dentin and Enamel Erosion
 - Dental Erosion and Diet
 - Saliva
 - The Salivary Pellicle
 - Gastroesophageal Reflex Disorder (GERD)
 - Patient Age and Risk Factors
- Caries versus Erosion
 - Key Differences Between Caries and Dental Erosion
 - The Role of Fluoride in Caries Reversal
 - Fluoride and Dental Erosion
- Current Methodologies to Assess Dental Erosion
 - Lesion Progression Methods
 - Surface Layer Measurements
 - Progression of the Erosive Lesion
- Clinical Strategies to Prevent and Manage Dental Erosion
 - Risk Assessment for ETW
 - Clinical Examination
 - The Basic Erosive Wear Examination (BEWE)
 - Challenges in the Management of ETW
 - Preventing and Managing Dental Erosion
 - Oral Hygiene Practices and Home Care
 - Professional Care
- Conclusions
- Course Test
- References / Additional Resources
- About the Authors

Overview

The dental research community has made great strides in preventive dentistry over the past several decades, with breakthroughs such as the introduction of fluoride and tartar control dentifrices, enhanced sensitivity reduction approaches and fluoride varnishes. In spite of these advances, dental erosion has become a major new challenge for dental professionals.

First identified approximately 25 years ago, the prevalence of dental erosion has increased dramatically in children, adolescents and adults ever since. This is of particular concern since the enamel and dentin loss associated with this multifactorial condition is irreversible.

Confusion exists, however, regarding the differences between dental erosion and dental caries. Although there are some similarities between these two unique processes, there are critical differences related to the etiological factors, the long-term effects, and the best ways to help manage these issues for each patient.

Over the past several years, we have learned a great deal about dental erosion. Fortunately, we have also learned about important strategies to help prevent it. This course provides you with the most current learnings on dental erosion, including research on the efficacy of fluoride sources to help prevent its initiation and progression.

Learning Objectives

Upon completion of this course, the dental professional should be able to:

- Discuss trends related to the prevalence of dental erosion.
- Understand similarities and differences between dental erosion and caries.
- Explain the etiology of dental erosion to patients.
- Discuss the long-term effects of dental erosion.
- Explain strategies for diagnosing and managing patients at risk for dental erosion.

Background

Dental erosion is a condition that results from an excessive exposure to erosive acids, either of extrinsic (dietary) or intrinsic (gastric) origin. First quantified on a wide scale basis in the United Kingdom,^{1,2} and also throughout Europe,³⁻⁷ this problem later gained significant interest on a more global scale.^{8,9} This condition is highly relevant to oral health professionals, and it presents these professionals with challenges regarding its treatment. From a patient's point of view, dental erosion can be associated with esthetic problems and pain

from dentin hypersensitivity. It can also impact long-term tooth function. From the oral health care professional's point of view, it can be very difficult to manage the condition; it sometimes requires changes in patient habits, which can present a significant hurdle.

In most cases, dental erosion does not present as a single condition. It is one part of a broader, multi-factorial 'umbrella' condition referred to as erosive tooth wear (ETW) (Figure 1). ETW is a growing problem, seen day to day in general practice (Figure 2). It includes different factors, including dental erosion, abfraction, abrasion and attrition; alone or in combination. Generally, ETW is classified according to the specific mechanism that is responsible for the wear. While the mechanism for tooth wear resulting from erosion is chemical, abfraction, abrasion and attrition are the result of physical forces.

In the past, particularly in the US, dental professionals often associated tooth wear with occlusion and bruxism. But the fact is it probably has more to do with acid. Changes on the lingual surfaces of eroded teeth, for example, are likely the result of a combination of acid and repetitive, frictional forces from the tongue.¹⁰ It is not from occlusion or any type of a bruxism-type movement. There are two distinct processes at work, which highlights the



Figure 1. Erosive Tooth Wear (ETW) is an umbrella term that includes dental erosion, abfraction, attrition and abrasion, alone or in combination.

complexity of the problem. Regardless of which forces are at play in an individual patient, the net clinical outcome is tooth surface loss.

- Dental erosion is an outcome resulting from the dissolution of dental hard tissue by either intrinsic or extrinsic acids that are not of biological origin.
- Abfraction is a form of physical wear along the gingival margin that is not caused by bacterial acid activity.
- Abrasion is a form of physical wear that is the result of mechanical interactions, such as tooth brushing or repetitive contact of a foreign object, with opposing tooth surfaces.
- Attrition is a form of physical wear that occurs as the result of one tooth coming into contact with another and is often associated with bruxism (tooth grinding).

Clinically, ETW is often associated with a combination of tooth wear processes, with dental erosion being the most common component. In addition, dental hygiene habits, such as brushing with a hard-bristled toothbrush or brushing too soon after taking in acid-containing food or beverages, can have an impact on tooth wear. Excessive tooth brushing can also remove significant portions of the acquired dental pellicle. Pellicle serves as a natural protection against both erosive acids and frictional wear. When teeth are brushed directly before eating or drinking, the thickness of the pellicle, and therefore its ability to protect exposed tooth surfaces, is reduced. Soon after brushing, the pellicle begins to be restored. Many dental professionals now suggest waiting for 1-2 hours after brushing before consuming acid-containing foods and beverages,¹¹ giving the pellicle sufficient time to regain a reasonable level of defense.

As we are all aware, people are living longer. If we were born today, our average life expectancy might be 100 years of age. Our medical colleagues recommend that we eat more fresh fruit and vegetables, which is sometimes a more erosive diet, in order to combat certain diseases, such as diabetes and cardiovascular disease. In addition, our consumption of acidic soft drinks is increasing dramatically year-on-year. Data comparing populations in both the



Figure 2. Severe erosion on a patient consuming 1.5 gallons of Kombucha fermented drink daily.

UK and US suggest we can anticipate finding a significant level of dental erosion in the general population,¹² with even higher numbers anticipated for specific high risk groups.^{2,4,13} The evidence suggests the presence of erosion is growing steadily.^{8,14} A recent study in Europe showed that 30% of young adults, 18-35 year olds, had dental erosion, much of which may be attributed to excessive consumption of soft drinks.¹⁵

'Baby Boomers' can represent a large patient population in many dental practices. These patients are much different than their similar age counterparts from 20-40 years ago. Patients now live longer, keep their teeth longer, and they're more physically active. Many of them are aesthetically sensitive, with different kinds of demands; they're not willing to settle for extractions and dentures. In fact, edentulism rates in the United States have decreased significantly in this age group, from about 45% in 1974 to around 17% now.¹⁶ This means that, for these patients in particular, we have far more teeth needing attention than we did several years ago. This needs to be acknowledged and properly managed.

It is not unreasonable to surmise that the increasing life expectancy, coupled with maintaining a healthier lifestyle involving a more acidic diet, may well lead to more and more cases of dental erosion. That is, of course, unless we put preventative measures in place to help address these concerns before significant damage is caused. As a start, dental professionals need to be far more proactive at looking for erosion, particularly at the earliest stages of the condition, and recommending the use of products that have been demonstrated to be effective at helping to prevent its initiation and progression.

Etiological Factors

Acid erosion involves a chemical process, a dissolution of hard tissue structures, without bacterial involvement. The erosive process is not necessarily a simple one. One way to think about it is by focusing on changes happening in the saliva. When the pH drops in the saliva, it then drops in the acquired, or salivary, pellicle. After that, acidic changes occur on the tooth surface, which initiates the series of events that lead to tooth surface loss. In reality, nothing happens on the tooth until it happens first in the saliva. As such, saliva has become a primary area of interest to monitor.

Poor salivary flow impacts clearance of acids and buffering, and therefore delays a return to the resting pH. The saliva also supplies the pellicle; pellicle helps prevent and stop progressive erosion unless overwhelmed by a strong acid challenge. Synergistic wear may occur, for example, by the tongue abrading softened enamel palatally and lingually.¹⁰

An excellent technical description of dental erosion has been offered by Ganss: "Dental erosion can be defined as dissolution of tooth by acids when the surrounding aqueous phase is undersaturated with respect to tooth mineral. When the acidic challenge is acting for long enough, a clinically visible defect occurs. On smooth surfaces, the original luster of the tooth dulls. Later, the convex areas flatten or shallow concavities become present which are mostly located coronal to the enamel-cementum junction. On the occlusal surfaces, cusps become rounded or cupped and edges of restorations appear to rise above the level of the adjacent tooth surfaces. In severe cases, the whole tooth morphologically disappears and the vertical crown height can be significantly reduced. The result of continuing acid exposure,




Figure 3. (a) SEM showing loss of enamel and, (b) at greater magnification, the softened layer at the advancing front of the lesion.

Images courtesy of Karger.¹⁸

however, is not only a clinically visible defect, but also a change in the physical properties of the remaining tooth surface. It is recognized that erosive demineralization results in a significant reduction in microhardness, making the softened surface more prone to mechanical impacts. Although independent in origin, erosion is linked to other forms of wear not only because it contributes to the individual overall rate of tooth tissue loss, but also by enhancing physical wear.¹⁷

In terms of acid erosion per se, we first have a surface softening, followed by surface loss, as depicted in Figure 3. Surface loss occurs as a result of frictional forces impacting the softened tooth mineral, followed by the initiation of a second softened layer, which is partly demineralized. The affected area of the tooth is susceptible to further frictional challenges, which leads to additional tooth surface loss.

Because of the interaction between frictional forces and acid, we have to consider tooth wear the result of a rather complicated process. In general, ETW is a multifactorial process that involves acid erosion and frictional forces of abrasion and attrition.

Differences Between Dentin and Enamel Erosion

One factor that complicates the issue even further is that the tissues involved in dental erosion, both enamel and dentin,

Table 1. Composition of enamel and dentin.

Component	Enamel	Dentin
Inorganic Material	85%	47%
Organic Material	2%	33%
Water	11%	21%

are very different (Table 1). While enamel is approximately 85% mineral, combined with a small amount of collagen, organic material and water, dentin is highly organic. Dentin is comprised of about 47% mineral, with the remainder a combination of organic matter and water. Due to the difference in makeup, dentin reacts much differently to erosive activity and wear than enamel (Figure 4).

One way to think about the dental erosion process is to consider how adhesive bonding procedures, using 37% phosphoric acid, condition the tooth surface to enhance retention of the material. After placement of the acid, the result is typically a visibly chalky appearance (Figure 5). The surface has been demineralized, or etched. That is the same thing that happens on a slightly different scale every time there's an acid challenge in someone's mouth, from whatever source. The primary




Figure 4. Tooth surfaces after an acid challenge. (a) Erosive demineralization of the enamel shows loss of surface between enamel prisms, with the prisms themselves remaining intact. These areas are susceptible to loss due to subsequent frictional forces. (b) Dentin surface subjected to the same erosive challenge reveals a surface that is much different from enamel. Demineralization of the intertubular matrix has occurred, and some of the dentinal tubules become opened, which can lead to sensitivity.

Images courtesy of Karger.¹⁸



Figure 5. Typical etching pattern on teeth after use of a 37% phosphoric acid treatment.

difference is that one challenge is controlled, and limited to a single exposure, while the other can occur over and over in the mouth, over a prolonged period of time. After an erosive acid challenge on enamel, for example, enamel prisms remain (Figure 4a). Once these demineralized prisms of enamel are present, they are highly susceptible to abrasive forces. Micron by micron, the tongue, food, occlusal forces, etc., will break these susceptible areas off and begin a repetitive cycle of tooth surface softening and loss.

Although the same factors are at play on dentin, the overall process of dental erosion on dentin is somewhat different. When dentin is attacked by erosive acids, the result is a demineralized organic matrix. The mineral

portion becomes highly demineralized (Figure 4b). This is very important, for example, in adhesive dentistry and when doing bonding type techniques. It is not hard to imagine how erosive acids predispose the dentin to surface loss and wear. These processes also expose open dentinal tubules, which can then lead to tooth sensitivity.

Another complicating factor is that dentin is subject to degradation by proteolytic enzymes, the MMPs, among other things.¹⁹ As we try to understand these processes, we have to include this as one of the risk factors in the degradation, or changes, that occur in dentin erosion. If the dentin becomes soft and liquefied, this will have a significant effect on the magnitude, the extent and the rapidity of tooth surface loss.

To briefly summarize, in the case of enamel erosion, we see more demineralization and bulk tissue loss, which is primarily due to the higher mineral content in enamel. Importantly, many of these changes occur at a pH of less than 4. Dentin, on the other hand, has overall less demineralization and bulk tissue loss under an acid challenge, has a softer matrix and is more susceptible to surface loss due to frictional forces. Changes in dentin typically occur at somewhat higher pH, usually above pH 4.

From a clinical standpoint, erosive processes can appear to be contradictory (Figure 6). In reviewing this figure, one might question how

there is so much dentin loss, while the enamel appears to be much less affected. Relating this image to the discussion above, it is likely that in this patient the erosive challenge wasn't at a very low pH. It may have been a higher pH, still under 5.5 or so, but at a pH that didn't have a significant impact on the enamel. Because it wasn't a low enough pH, there was a much greater erosive effect on dentin than enamel. At the same time, the area identified by the green arrow shows a much different situation. This area likely involves actual frictional forces from the opposing teeth, which has resulted in more enamel loss. The question becomes: how do you have, in the same mouth, this level of discrepancy? This dilemma serves to highlight the complexity and the clinical challenges that dental professionals face in dealing with this problem. One answer is to make sure that dental professionals are trained to assess this condition from multiple perspectives.

One problem, from an epidemiologic standpoint, is that dental erosion is often not noted on a patient's chart, especially in the United States. This contrasts significantly with Europe, Australia and some South American countries, where the assessment of dental erosion has become a routine practice. In the past, unless there was pain or some type of cosmetic problem, patients did not seek treatment for erosion, and most dentists didn't offer care for it. As dental erosion has become more of an issue, it is hoped that all dental practitioners develop a greater awareness of the problem and become more equipped to



Figure 6. Image shows high level of ETW on the dentin surfaces, yet leaves some areas of enamel less affected.

Photo courtesy of Dr. Michael Nelson.

help manage their patients that are either at risk, or are already experiencing some level, of dental erosion.

Dental Erosion and Diet

Most researchers consider acid-containing soft drinks and beverages as primary culprits in the growing incidence of dental erosion.^{8,9,14,15,20-22} Over the past several decades, serving sizes in the US have increased dramatically.^{8,23} The average drink size, in the 1950s, was slightly less than seven (7) fluid (fl.) ounces (207 mL). By the 1960s, this average serving size had increased to twelve (12) fl. ounces (355 mL), and by the late 1990s had increased still further to twenty (20) fl. ounces (532 mL) (Figure 7). In many restaurants, the largest sizes that are sold often contain 42-44 fl. ounces (1,242-1,301 mL), and free refills are commonly available. Between 56% and 85% of children at school have been reported to consume at least one (1) soft drink daily, with 20% consuming an average of four (4) or more servings every day.^{8,24}

In the US, current caloric intake has been shown to be up to about 4,000 calories per day per person. Approximately 25% of this caloric consumption is in the form of sugar-sweetened beverages or, at the very least, added sugars to the diet. A study by Credit Suisse compared the gross domestic product of various nations



Figure 7. Beverage sizes have grown significantly. In the 1950s, the average cola drink in a US restaurant was 7 fl. ounces (207 mL). Serving sizes have increased steadily ever since; today, the average serving size is over 30 fl. ounces (887 mL), with individual "large" portions often providing 42-44 fl. ounces (1,242-1,301 mL), or more, of beverage.

Annual Soda Consumption versus GDP (Per Capita)




Figure 8. Annual global consumption of sodas vs. GDP per capita clearly demonstrate a much higher level of consumption in the US compared to other nations. Adapted from Sugar, Consumption at a Crossroads. Credit Suisse.²⁵

versus their annual consumption of sugar-sweetened beverages.²⁵ As shown in Figure 8, the US is well above other countries in this assessment, which indicates a very high risk for dental erosion in this country, due to the low pH and high acid levels in sugar-sweetened beverages. In the US, the average person consumes over 40 gallons (151 L) of sugar-sweetened beverages per year.^{25,26} It is no surprise that the US has obesity problems, caries problems, and other related side effects.

The severity of erosive acid attacks depends on multiple factors, such as the pH (Table 2), the titratable acidity and the buffering capacity of both the beverage, or food, and the saliva of the individual ingesting the beverage. Another primary factor is the contact time of the acid on the teeth. The longer the teeth are subject to an erosive acid challenge, the more likely they will be to undergo erosive changes.

In addition, all acids are not alike with regard to their erosive potential. Studies have shown that citric and lactic acids have a higher erosive potential than acetic, maleic, phosphoric and

tartaric acids, although all of these dietary acids have some degree of erosive potential. When included in products containing other ingredients, such as calcium, phosphate and/or fluoride, the erosive potential of an acid can be significantly decreased. For example, acidic beverages, when supplemented with calcium, phosphate and fluoride, have been shown to have a reduced erosive potential compared to controls.²⁷ In addition, yogurt, which has a relatively low pH, has little erosive potential due to its high calcium and phosphate content.²⁸

Consumption of a single acidic beverage and drinking it normally has little impact on dental erosion. Although the pH of saliva does drop while drinking it, the saliva will generally provide sufficient buffering to quickly re-establish a neutral pH. If people swish their drinks, or if they sip these beverages over long periods of time, there is a much higher likelihood of having problems. One way to help minimize the potential for prolonged contact of acidic beverages with the teeth is to drink through a straw, as this may help to minimize contact with the teeth.²⁹

Table 2. pH values of common beverages.

Beverage	pH
Pure Water	7.0
Milk	6.6
Tea	6.2
Coffee and milk	5.3
Black Coffee	5
Tomato juice	3.7
Sprite	3.29
Diet Pepsi	3.03
Gatorade	2.92
Dr. Pepper	2.90
Fruit Punch	2.82
Orange Juice (Minute Maid)	2.64
Coke Classic	2.53
Reference: Battery Acid	1.0

Adapted from Jain et al., General Dentistry.³⁰

While there is growing awareness of dietary issues related to the high consumption of soft drinks, particularly among school aged children, the issue of dental erosion is not limited to consumption of only these beverages. Equally challenging to the enamel surface is excess exposure to fruits, acid containing vegetables, fruit juices, wine, and other dietary foods and beverages that are

otherwise considered healthy alternatives to soft drinks. It is difficult to protect against an increasing erosive challenge, no matter what kind of diet is consumed.^{28,29,31-33}

One of the driving factors for dental erosion is the duration of contact between dietary acids and the teeth. As a result, there is some perception that high viscosity drinks may be more erosive than those with lower viscosity, due to their tendency to be retained in the mouth for longer periods of time. Recent in vitro studies, however, have suggested that the opposite might be true,^{34,35} possibly a result of the ingredients that enhance viscosity actually being able to slow the release of acids from the beverage, thus causing less damage. This area of research will be interesting to watch, as it will take more robust models and clinical trials to confirm whether increased viscosity enhances erosion, or helps to prevent it.

Saliva

One of the most important factors influencing the progression of dental erosion is saliva, especially in our age of polypharma. Even before an acid attack takes place, saliva flow is often increased as a response to stimuli such as smell or chewing. A high saliva flow rate helps increase buffering, dilution and clearance of acids from the mouth, which is extremely important during an erosive acid challenge. A low salivary flow rate can have a number of negative effects. Saliva flow can be inhibited as a side effect to numerous medications, both prescription and over-the-counter. Medical conditions, such as xerostomia, dehydration, and salivary gland dysfunction can all put teeth at risk for erosion.³⁶

The average adult in the US, over the age of 65, takes six or more prescription medications. Many of these medications have the potential to adversely impact saliva, or salivary flow rates, which can increase the risk of dental erosion. Low levels of saliva can impact the rate of pellicle maturation, and could also result in elevated risk of erosion due to excessive tooth brushing, chewing hard foods and bruxism. At the very least, changes in the saliva can make the process of dental erosion and toothwear much more complex.³⁷

In older populations, where there is a combination of hyposalivation and polypharma, this can be a significant problem. These individuals can be exposed to very low pH acids in their mouth. When people are asleep at night, they only have about one-tenth of the saliva they have during the day. If someone is over 65, they likely have only half of the saliva they had when they were 20. Without sufficient saliva, it is difficult to provide sufficient buffering and clearance of acids, particularly if the patient suffers from acid reflux.

The chemical composition and buffering capacity of saliva are also important factors. If saliva has a high concentration of bicarbonate, it has an increased capability of neutralizing and buffering erosive acids, and if supersaturated with respect to calcium and phosphate, it is better equipped to reverse low levels of initial softening that might occur. Saliva that is undersaturated with respect to calcium and phosphate has little ability to help protect tooth surfaces against erosive acid challenges.³⁸

The Salivary Pellicle

Besides saliva, another key biological factor that can influence the progression of dental erosion is the pellicle. The salivary, or acquired, pellicle is a protein-based layer that covers all orally exposed surfaces of the teeth. Although this pellicle film can be modified somewhat through brushing, the pellicle film is essentially never removed from the teeth, with the exception of during a dental prophylaxis (prophy). During a prophy, the abrasiveness of prophy pastes is sufficiently high to enable the complete removal of this film from the tooth surface, exposing the natural enamel surface. The reason for removing the pellicle during the prophy is to allow for the complete removal of extrinsic stains from the tooth surface. While chemical cleaning agents like peroxide are able to bleach stains from an intact pellicle, prophy pastes actually remove the pellicle. In order to restore its natural level of protection, the saliva generates a new, fresh pellicle within just a few short hours. Though thin in absolute terms, the pellicle provides protection to the tooth surface against extrinsic acid damage.^{8,36,39}

When acidic foods and beverages are taken in excess, the pellicle layer can be overwhelmed by either the sheer volume of dietary or gastric acids, the high titratable acidity of a particular beverage, or a complex combination of factors such as the mineral content of saliva.³⁸ When this occurs, the surface enamel softens quickly, and even the fluoride rich outer enamel, which provides a measurable level of ‘second defense’ against cariogenic acids, cannot defend itself against a strong erosive acid challenge. The outer enamel becomes softened and susceptible to damage; due primarily to a multitude of potential “tooth wear” factors present in the mouth.¹⁴

Gastroesophageal Reflex Disorder (GERD)

Another issue that has a significant impact on dental erosion is gastroesophageal reflux disorder (GERD). Significant increases in severe erosion are seen in the older population, and between 10 to 20% experience gastric acid symptoms.⁴⁰ In 2015, over 113 million prescriptions for antacid medications were written, representing over \$13 billion in sales.⁴¹ Even more worrisome is silent GERD. A recent study found over one-third of patients may have silent GERD, where the reflux actually gets up into their mouth but the patients are asymptomatic.⁴² This can have a significant impact on dental erosion. Controlling GERD has been shown to have a positive impact on dental erosion. In one study, patients who had dental erosions that were recorded as active lesions were prescribed proton pump inhibitors (PPIs) to help manage their gastric reflux issues. In 86% of cases, progression of the erosive lesions was stopped.⁴³ These results demonstrated that by suppressing the gastric acids, the incidence of erosion was reduced; and the progression of dental erosion was effectively managed.

Patient Age and Risk Factors

ETW is episodic throughout life and can start in infancy. It is generally accepted that deciduous ETW may be indicative of future erosion problems in the permanent dentition. Although studies in the literature have suggested the prevalence of dental erosion to be somewhere between 7-74%, an overall ETW prevalence of 30% has been found in a meta-analysis of




Figure 9. Multifactorial ETW with erosion and attrition.



Figure 10. The presence of NCCL lesions suggests the likelihood of improper brushing habits.

studies in teenagers and young adults with at least one erosive lesion, and the condition becomes even more prevalent with age.⁴⁴ Different etiologies typically play more of a role at different ages. Figure 9 shows a case involving erosion and attrition.

Risk factors for ETW include: 1) dietary habits (amounts, frequency, manner of consuming acidic foods and beverages); 2) gastric reflux (GERD, bulimia, pregnancy vomiting); 3) xerostomia - reduced salivary flow decreases acid dilution and clearance; and, 4) exposure to mechanical insults such as hard foods, improper toothbrushing and bruxism. The severity of acid attacks varies with the pH of acid and its buffering capacity, whether a drink is swished/sipped/gulped or taken with a straw, and its contact time, the thickness of the acquired pellicle and salivation. Reduced

salivary flow represents the greatest risk factor and must be evaluated. ETW is irreversible and, if observed in children or adolescents, it can be expected to progress unless intervention occurs.

In looking at minor erosion, the patient's age, habits, and whether wear is physiological or pathological should be considered. Early diagnosis is especially important as patients typically do not seek care until they experience pain or an esthetic problem.

Conducting an evaluation for dental erosion provides dental practitioners with a window into some of the other habits of a patient. For example, recession above the NCCL lesions in Figure 10 suggests that this patient has likely been brushing with a stiff toothbrush and abrading the tissue away; this has resulted in minimally attached gingiva, recession and a very deep erosive lesion. Care must be taken to appropriately manage this type of condition, in addition to the other issues that patient is experiencing.

Caries versus Erosion

We know that caries can occur on any tooth surface. However, it is generally accepted that caries occurs under plaque and is the direct result of bacterial acids. The primary acid that causes caries is lactic acid, a byproduct of the breakdown of fermentable carbohydrates (primarily sugar) by plaque bacteria. While the most dominant bacteria responsible for caries are *S. mutans*, other bacteria, such as *lactobacillus*, have also been suggested as contributors to various aspects of the caries process.^{45,46}

Erosion, on the other hand, is a result of the direct action of extrinsic, dietary acids; such as those found in carbonated drinks and fruit juices or intrinsic acids, such as from GERD. Dietary acids include phosphoric, citric, and other acids commonly used to impart the tart, tangy flavors we associate with acidic foods and beverages. Although "diet" drinks are generally "sugar free," and thus more acceptable from a caries standpoint, the acid content of the diet beverages is no different from their sugar-containing counterparts. From

the standpoint of acid content, “sugar free” drinks offer no advantage when it comes to their potential to cause dental erosion.

Key Differences Between Caries and Dental Erosion

Generalities can be confusing. Caries is often described as the loss of minerals by the direct action of acids on the teeth, and dental erosion is also defined in a similar way. While both statements are true, of primary importance is the type of acid, where the acid comes from and specific sites on the tooth surface to which these acids are directed. It is important to differentiate enamel damage due to caries vs. damage that results from dental erosion. Both the etiology and symptoms of these two processes differ significantly, as do the appropriate management strategies for each (Table 3).

One major difference between caries and dental erosion needs to be clearly understood. Caries is a process that begins with demineralization and, at early stages, can be reversed, either through the natural process of remineralization or through enhanced remineralization due to fluoride therapy. Dental erosion, on the other hand, is essentially a non-reversible process that results in permanent damage to the tooth structure.

The Role of Fluoride in Caries Reversal

Dental caries is an infectious disease caused by the complex interaction of cariogenic (caries-causing) bacteria with carbohydrates (i.e., sugars) on the tooth surface over time. Cariogenic bacteria metabolize carbohydrates for energy and produce organic acids as byproducts. The acids lower the pH in the plaque biofilm.⁴⁷

The hydroxyapatite of tooth enamel is primarily composed of phosphate ions (PO_4^{3-}) and calcium ions (Ca^{2+}). Under normal conditions, there is a stable equilibrium between the calcium and phosphate ions in saliva and the crystalline hydroxyapatite that comprises 96% of tooth enamel. When the pH drops below a critical level (5.5 for enamel, and 6.2 for dentin), it causes the dissolution of tooth mineral (hydroxyapatite) in a process called

demineralization. When the natural buffer capacity of saliva elevates pH, minerals are reincorporated into the tooth through the process of remineralization.⁴⁷




The initial stage of the caries process results in white spot formation, a result of acid penetration and solubilization of some (but not all) of the subsurface mineral (Figure 11A). Left untreated, this subsurface damage can progress to a point where the crystal can no longer provide sufficient support to the enamel surface structure, and the surface collapses (cavitates).

The caries process can be affected in several ways. One of the most effective methods to prevent caries is by promoting remineralization and slowing down demineralization. This can be accomplished with fluoride therapy. It is widely accepted that the regular use of fluoride, such as in dentifrice and drinking water, is extremely effective at preventing dental caries. In 1999, the US Center for Disease Control (CDC) issued a statement that water fluoridation is one of the 10 most important public health measures of the 20th century.⁴⁸ Fluoride's presence in low concentration and high frequency is more effective at preventing caries than high levels of fluoride used in low frequency. Because water fluoridation is not available in many countries, dentifrice is considered to be one of the most important sources of fluoride globally.

When fluoride is present in oral fluids (i.e., saliva), fluorapatite, rather than hydroxyapatite, forms during the remineralization process. Fluoride ions (F^-) replace hydroxyl groups (OH^-) in the formation of the apatite crystal lattice (Figure 11B), resulting in a stronger, fluoridated tooth mineral (fluorapatite). Fluorapatite is less soluble than hydroxyapatite, even under acidic conditions. Because fluorapatite is less soluble than hydroxyapatite, it is also more resistant to subsequent demineralization when acid challenged.

Caries is generally considered to be a subsurface phenomenon. With fluoride treatment, a noncavitated lesion can be remineralized with fluorapatite and have greater

Table 3. Key differences between caries and dental erosion.

		
Key Comparisons	Cavities	Erosion
Type of process	Mineral Change	Mineral Loss
Tooth site(s) affected	Enamel & dentin	Enamel & dentin
Primary cause	Bacterial acids	Dietary acids
Primary site(s) of damage	Subsurface, under plaque	Exposed, plaque free surface
Conditions	Exposure to weak acids for prolonged periods of time, usually at a pH above 4.0	Repeated exposure to dietary or gastric acids, generally below pH 4.0, for short time periods
Result	Sub-surface phenomenon with intact outer layer of enamel	Surface softening leading to loss of surface mineral
Reversible?	Reversible in early stages	Irreversible surface loss
Contributing factors	Buffering by saliva helps neutralize bacterial acids	Saliva and pellicle overwhelmed by dietary and gastric acids
Preferred therapeutic approach	Prevention as well as reversal of early damage	Prevention is critical for managing
Fluoride effectiveness	-----	-----
Sodium fluoride	Yes	Minimal
Sodium monofluorophosphate	Yes	Minimal
Stannous fluoride	Yes	Yes




Figure 11. A) Demineralization – the caries formation process. Damage occurs in subsurface regions of the enamel, leaving an intact outer layer on the enamel surface. B) Remineralization – the caries reversal process. Caries is reversed through the process of remineralization, in which calcium, phosphate and fluoride are incorporated in the areas damaged due to demineralization processes, resulting in a stronger, fluoridated mineral.

resistance to subsequent demineralization than hydroxyapatite. Even at very low concentrations, fluoride is effective as an anticaries agent.⁴⁹

In the US, there are three commonly used sources of fluoride in oral care products; sodium fluoride (NaF), sodium monofluorophosphate (SMFP) and stannous fluoride (SnF₂). All three of these fluoride sources provide the important F⁻ ion, which

both inhibits demineralization and promotes remineralization of damaged tooth mineral. In addition, SnF₂ is considered to have unique properties, as it provides efficacy against bacterial acids in addition to its fluoridating benefits.

Fluoride and Dental Erosion

While dental erosion, like caries, is a mineral process, the erosion process follows a somewhat different pathway.^{38,50} There is little




Figure 12. Dental Erosion. Erosion processes result in permanent loss of surface mineral structure.




Figure 13. Stannous fluoride (SnF_2) is unique among the sources of fluoride routinely used for toothpaste formulations.

possibility of reversal, as erosive acids are able to overwhelm the protective pellicle layer and soften outer surfaces of the tooth; these softened surfaces can then be lost to abrasive forces, resulting in permanent and irreversible loss of tooth structure (Figure 12).

It is well accepted that fluoride helps keep teeth strong. However, recent studies have demonstrated that all fluorides are not alike with their ability to help prevent dental erosion. While there would likely be a greater incidence of dental erosion in the absence of fluoride, the data suggest that most fluorides do not provide a high level of benefit against the increasing levels of challenge teeth are facing in today's environment. In spite of the fact that almost 100% of the world's toothpastes contain fluoride, the incidence and prevalence of dental erosion both appear to be on the rise. These data suggest many fluoride products

may not be sufficiently effective to protect teeth against erosive acid challenges. However, one of the currently used sources of fluoride, stannous fluoride (SnF_2) (Figure 13), has been demonstrated in a broad range of studies to be unique in its ability to help prevent the initiation and progression of dental erosion. These include both laboratory⁵¹⁻⁵⁴ and human in situ erosion clinical studies.⁵⁵⁻⁶⁰ Different from other sources of fluoride used, stannous fluoride deposits a retentive, acid resistant barrier layer onto exposed tooth surfaces that is protective against both the initiation and progression of dental erosion (Figure 14).⁶¹

Current Methodologies to Assess Dental Erosion

Methodologies used to assess dental erosion efficacy fall into 2 groups - those measuring total mineral loss from enamel (and dentin) due to lesion progression, and those measuring the surface properties of erosive lesions.

Lesion Progression Methods

Lesion progression can be assessed using sound enamel slabs subjected to cycles of acid challenges and salivary remineralization in vitro, or by in situ testing with patients drinking beverages that deliver acid challenges to enamel slabs worn in an appliance (Figure 15). Using the same methods, the efficacy of preventive measures to inhibit lesion progression can be assessed. Lesion progression in enamel and dentin can be measured using microradiography, contact profilometry, and non-contact (optical)




Figure 14. SnF₂ helps prevent erosion by depositing an acid resistant barrier layer on exposed tooth surfaces, thus preventing the initiation and progression of irreversible damage.
Adapted from Faller; Cosmetics & Toiletries, 2012.⁶²



Figure 15. Human enamel specimens mounted into appliances worn for human *in situ* erosion prevention studies.

profilometry; the latter two additionally measure surface roughness. A fourth option, confocal laser scanning microscopy (CLSM), measures total lesion progression and the softened zone of mineral caused by an erosive acid. A recent study compared CLSM, contact profilometry and non-contact profilometry and found all three methods gave similar results in measuring the loss of enamel.⁶³

Surface Layer Measurements

Surface layers can be assessed using *in vitro* or *in situ* models. Methods of analysis can include scanning electron microscopy (SEM) (Figure 16),

quantitative light fluorescence or optical coherence tomography which measure changes in the surface zone and are used to determine surface roughness. Additional methods include acid solubility testing and atomic force microscopy (Figure 17). Replica SEMs can be used with *in situ* research to measure changes in the surfaces of eroded lesions or softened zones of enamel lesions over time. An additional method of analyzing the surface layers, secondary ion mass spectroscopy (SIMS), can be used to measure the composition of the surface layer and to show material firmly deposited at the surface.




Figure 16. SEM showing loss of enamel (a) and, at greater magnification, the softened layer at the advancing front of the lesion (b). Images courtesy of Karger.⁶²




Figure 17. Atomic force micrographs show the decreased loss of structure on the enamel slab treated with stabilized stannous fluoride toothpaste. Stannous fluoride deposits are insoluble in concentrated acid and provide enhanced protection, especially at low pH.

Hardness recovery, hardness loss inhibition, surface composition/fluoridation and solubility reduction measurements can all be used to assess the mechanisms and efficacy of a proposed preventive therapy against dental erosion. Protocols used include surface preparation using an acid challenge and measurement, treatment steps, surface measurements after secondary acid challenges (typically citric acid, or for *in situ* studies orange juice), and then measuring specimen hardness again after re-immersion in saliva.^{51-55,58,59}

Progression of the Erosive Lesion

The 3 stages in progression of an erosive lesion are initial surface softening, progressive loss of enamel and the creation of a lesion that involves dentin. By focusing on lesion

progression models, the ability of a given therapy to protect against erosive progression can be determined. Two ways to help prevent dental erosion are to protect the tooth surfaces by, in effect, 'galvanizing' the surface with deposits that are acid-resistant, and/or to remineralize the surface between acid challenges. Superior efficacy for SnF₂ in reducing surface loss when measured *in vitro*, using microradiography (Figure 18) following several cycles of erosive challenges, immersion in saliva, and treatment with fluorides (Figures 19, Table 4) has been demonstrated in multiple studies.⁵¹⁻⁵⁴

Reduced progression of erosive lesions has also been observed with stabilized SnF₂ toothpaste *in situ* compared with sodium




Figure 18. Microradiography is used to quantitatively measure tooth surface loss in *in vitro* lesion progression models.

Adapted from: Eversole et al, 2014.⁵¹




Figure 19. Surface enamel loss with marketed dentifrices using a lesion progression model with microradiography.

Adapted from: Faller et al, 2011.⁵³

fluoride toothpaste. In a study by Hooper and colleagues, the benefit of stannous fluoride increased over time (Figure 20).⁵⁵

In a study by West et al (Figure 21), highly significant erosion protection benefits were found when comparing a stabilized SnF₂ dentifrice to a marketed dentifrice containing SMFP as the fluoride active and arginine bicarbonate.⁵⁹

As noted earlier, many foods and beverages have a pH below 4.0 and are highly acidic (Table 2). Although pH values are not an absolute predictor of erosive potential, it is important to know the general pH values for different foods and beverages. It is also important to consider these values when performing erosion testing, as different pH values will result in different results. During an erosive acid challenge, the tooth starts to

Table 4. Results and statistical analysis: Percent reduction in enamel surface loss comparing marketed dentifrices using a lesion progression model with microradiography.

Type of Formula Tested	Mean Surface Loss ± SEM (µm)	% Reduction vs. Control ^b
Stabilized SnF ₂ ^a	5.5 (1.2)	70.5
Mixed-active cavity protection ^c	16.0 (2.0)	14.0
SMFP/arginine bicarbonate ^d	17.1 (1.1)	8.2
Single active cavity protection ^b	18.7 (0.9)	-----

^a Crest® Pro-Health, The Procter & Gamble Company, Cincinnati, OH, USA

^b Crest® Cavity Protection, The Procter & Gamble Company, Cincinnati, OH, USA

^c Colgate® Cavity Protection, Colgate-Palmolive (UK) Limited

^d Colgate® Sensitive Pro-Relief™, Colgate-Palmolive (UK) Limited

* Mean ± SEM from Least Significant Difference Analysis

Means within the same bracket were not statistically significantly different at the 0.05 level of significance.




Figure 20. Superior erosion protection was demonstrated for SnF₂ in an *in situ* human clinical study.

Adapted from Hooper et al, 2007.⁵⁵

dissolve, in an effort to restore an equilibrium, by releasing calcium, phosphate and fluoride salts. At pH4, more than 90% of the fluoride salt released is present as fluoride ions, which protects the surface against demineralization.

At a pH of 2, it is overwhelmingly hydrofluoric acid that is present, rather than fluoride ions; as a result, insufficient fluoride ions are present to protect the tooth surface, which leads to dissolution of the surface layer of the tooth.




Figure 21. Results from a human *in situ* clinical lesion progression study comparing a stabilized SnF₂ dentifrice versus a marketed dentifrice formulated with SMFP and arginine bicarbonate.⁵⁹

Stannous fluoride has been shown to provide unique protection against acids, particularly at a low pH (e.g., orange juice, ~2.6).^{55,58,59}

There are a variety of analytical methods, models and protocols that can be used to study erosion. Some measure lost mineral while others measure changes in the surface zone. Protocols differ depending on whether studies will be used to analyze the surface zone or to produce erosive lesions and measure effects on the progression of erosion. Depending on the method used, dramatic differences can be seen in the observed efficacy of various topical agents. Importantly, SnF₂ dentifrices, in particular stabilized SnF₂ dentifrices, have been demonstrated, using a wide range of both *in vitro* and *in situ* human clinical studies, to provide significantly greater erosion protection than other fluoride sources.⁵¹⁻⁶⁰

Clinical Strategies to Prevent and Manage Dental Erosion

Pathological tooth wear was rarely seen in ancient civilizations, and the majority of it was



Figure 22. Facial view of dentition of a 23-year-old male with ETW, a condition that can lead to poor aesthetics, sensitivity, loss of function, and sometimes loss of self-esteem.

abrasion or attrition. While its prevalence and severity have increased in children and adults, ETW is a totally preventable condition for most individuals. Progressive ETW can lead to poor aesthetics, sensitivity, loss of function, and sometimes loss of self-esteem (Figure 22).

Early identification and prevention is key. By the time lesions are clearly visible to the patient, restorative intervention and life-long dental treatment may be required. Regularly

screening all patients for ETW makes it possible to diagnose erosive lesions at the earliest possible stage and allows implementation of preventive and treatment measures to preserve tooth structure and help stop further damage.

Risk Assessment for ETW

The risk assessment should consider all risk factors and include asking patients if they are aware of any tooth wear, or have sensitive teeth. The risk assessment includes determining dietary habits; oral hygiene regimens, the presence of gastroesophageal reflux (GERD) symptoms, eating disorders, xerostomia, and whether the patient is a lactovegetarian, taking acidic medications, or occupationally at risk (e.g., wine tasters, professional swimmers). A number of factors influence the rate of progression and extent of erosive lesions and must be considered during the risk assessment.

These factors (Table 5) include the frequency, amount and duration of exposure to erosive acid challenges, such as a high consumption of carbonated drinks, acidic fruits and vegetables, drinking alcohol, the manner in which acidic agents are consumer, e.g., holding or swishing acidic drinks in the mouth which prolongs contact; and, having acidic drinks at night when salivary flow is low.⁶⁴⁻⁶⁷

Clinical Examination

A full examination and tooth indexing for erosion, known as the Basic Erosive Wear Examination (BEWE), should be performed. The patient should be assessed for erosion, abrasion, attrition, abfraction, tooth wear etiology, recession, dentinal hypersensitivity, occlusion, salivary flow rate, and staining (which would suggest erosion is likely not occurring at that time) (Figure 23).

Table 5. Key risk factors influence the rate of progression and the extent of erosive lesions.



Key Risk Factors Influencing the Rate of Progression and Extent of Erosive Lesions	
Frequency of consuming erosive acid containing beverages	
Duration of erosive acid exposure	

Table 5. Key risk factors influence the rate of progression and the extent of erosive lesions (Continued).





<p>High consumption of acidic fruits and vegetables</p>	
<p>Drinking alcohol</p>	
<p>The manner in which acidic agents are consumed</p>	
<p>Having acidic drinks at night when saliva flow is low</p>	



Figure 23. Image of young 30-year-old presenting with multiple types of wear.





The Basic Erosive Wear Examination (BEWE)

The BEWE, introduced by Bartlett, Ganss and Lussi in 2008,⁶⁸ is used to assess the level of erosion. For this examination, the mouth is divided into six distinct areas for evaluation. Table 6 shows the criteria for sextant scores from 0 to 3, which are summed to obtain a cumulative score that is the basis for determining interventions (Table 7). The BEWE is a simple, quick index for screening a patient's erosion status.

Challenges in the Management of ETW

Management challenges for ETW include early diagnosis of erosive lesions, the initiation of preventive strategies and behavioral changes, and early intervention with minimally-invasive restorative procedures. Early diagnosis should include charting of erosive lesions, sensitive teeth, staining, and making note of areas of exposed dentin. In addition, a risk assessment should be performed.

Table 6. BEWE Scores and Criteria.

Score	Criteria	Image
0	No ETW	
1	Initial loss of surface texture	
2 ^a	Distinct defect; hard tissue loss involving <50% of the surface area	
3 ^a	Hard tissue loss involving ≥50% of the surface area	

^a Dentin often involved.

Table 7. Cumulative BEWE Scores and Management Guidelines.

Cumulative Score	General guidelines for management
No ETW: 0 - 2	<ul style="list-style-type: none"> • Routine maintenance and observation • Repeat at 3-year intervals
Low: 3 - 8	<ul style="list-style-type: none"> • Oral hygiene, dietary assessment • Routine maintenance and observation • Repeat at 2-year intervals
Medium: 9 -13	<ul style="list-style-type: none"> • Oral hygiene, dietary assessment • Routine maintenance • Fluoride measures • Avoid restorations • Repeat at 6-12 month intervals
High: ≥14	<ul style="list-style-type: none"> • Oral hygiene, dietary assessment • Routine maintenance • Fluoride measures • Repeat at 6-12 month intervals • Consider restorations

Patients should be taught preventive habits that reduce the risk of ETW. These include:

- Staying hydrated.
- Rinsing with water before brushing.
- Brushing with a stannous fluoride toothpaste.
- Not brushing for at least 1-2 hours after an acid challenge.

The bottom line in the management of ETW is early diagnosis, initiation of preventive

measures, and early intervention to avoid the need for extensive and invasive care.

Preventing and Managing Dental Erosion

Once a diagnosis of dental erosion is made, an overall preventive management program is needed. ETW management focuses on oral hygiene practices, home care, professional care, and individually tailored advice, depending on the level of severity found, to prevent further erosion and to manage the condition effectively.

There are a number of suggestions that can be made to all patients, and particularly to those in some of the higher risk categories. These categories are best defined as those individuals with a high consumption of dietary acids, such as colas (diet or regular are no different), fruit juices, wine, acidic fruits such as oranges, grapefruit, berries, apples, acidic vegetables such as rhubarb, tomatoes, any vegetables processed in vinegar such as canned beets, pickles, sauerkraut, and a host of other dietary components.

From the standpoint of Management Strategies, there are two main approaches. One is directed to the patient, while the other is geared toward the Dental Professional.

Oral Hygiene Practices and Home Care

It is recommended that patients with ETW brush for no more than 2 minutes and a maximum of twice-daily. Normally, manual and powered brushing cause virtually no enamel loss and minimal dentin loss. Some studies have suggested that power brushes, due to their ability to control the force of brushing, may be preferred over manual brushes.^{69,70} A common question is with respect to when is the best time to brush; before or after an erosive acid challenge. In a recent in situ study using enamel slabs, it took about 2 hours after an acid challenge before the enamel surfaces began to recover;⁷¹ this suggests that it may be wise to wait for at least 2 hours after an erosive acid challenge before brushing. If brushing before an acid exposure, make sure to use a product that provides an acid resistant barrier layer, such as a SnF₂ toothpaste, to protect the teeth against erosive acid attack. Without that barrier layer, the teeth could be susceptible to erosive softening, due to the effect of the brushing on pellicle thickness. Not brushing at all, of course, leads to other issues.

Patients with erosion can benefit from twice-daily use of SnF₂ toothpaste, because polyvalent metal ions interact with the tooth surface to form an acid-resistant insoluble layer. In fact, the recent consensus report by the European Federation of Conservative Dentistry notes that oral hygiene products containing stannous fluoride or stannous

chloride, such as toothpastes or mouth rinses, have the potential to slow the progression of ETW.⁶⁶ Additional options include recommending a calcium phosphate-based or bioactive glass home use product to promote remineralization.

Professional Care

Oral health education and advice must be individualized. In addition to recommending the use of stannous fluoride dentifrices, preventive care can include fluoride varnishes. Tooth surface protective coatings may also be indicated and dentinal hypersensitivity requires treatment or use of a desensitizing toothpaste (e.g., stannous fluoride). If intrinsic acid erosion is present (e.g., from GERD or bulimia), the patient should be referred for appropriate medical assessment and care. Restorative and bonding materials are only used if absolutely necessary to reduce sensitivity, improve esthetic considerations or restore function. For noncarious cervical lesions (NCCLs), it may be preferable to avoid restorative care, particularly for early lesions (Figure 10).

Patients should be given advice on simple, practical ways to reduce the risk of erosive tooth wear such as dietary advice and modifying habits. Table 8 contains a list of areas to consider when tailoring patient advice. Regular reassessment and monitoring are needed to determine if ETW has been halted and to provide patients with advice and care.

From the patient standpoint, probably the biggest area of concern is for those individuals with a high intake of acidic beverages. For those individuals, the recommendations are rather straightforward (Table 9). The first recommendation is to try to minimize the excessive intake of these beverages, as the pellicle can certainly accommodate some level of intake without issue. When consuming these types of beverages, it is generally considered best to drink with a straw, as this helps direct the acids past the teeth and into the mouth directly. Recommend drinking acidic beverages in a short period of time in order to minimize the overall time of contact of the acids with the enamel surfaces. Rinsing with water after drinking an acidic beverage will

Table 8. Individually tailored advice for patients.

Dietary Considerations for Sharing with Patients.
Reduce intake and frequency of acidic foods and drinks.
Eat cheese/milk after acid challenges.
Additional calcium content in food and beverages.
Drink cold rather than warm beverages (if acidic).
Rinse with sodium bicarbonate to help increase the pH after dietary acid challenges.
Chew nonacidic gum to stimulate saliva. Mint-flavored, sugar free gum should be suggested rather than citrus flavored gums, which are likely to be more acidic.

Table 9. Awareness for patients who ingest excessive amounts of erosive, acid containing beverages.

Erosion Awareness for Patients with High Intake of Acidic Beverages.	
Avoid excessive use of acid beverages.	
Use a straw.	

Table 9. Awareness for patients who ingest excessive amounts of erosive, acid containing beverages (Continued).




<p>Drink acidic beverages in a short period of time.</p>	
<p>Rinse with water after drinking to minimize exposure.</p>	
<p>Cold beverages are probably better than warm ones.</p>	
<p>Wait 1-2 hours after ingesting acidic products before brushing.</p>	

Table 9. Awareness for patients who ingest excessive amounts of erosive, acid containing beverages (Continued).



help dilute acids and rinse them away from susceptible tooth surfaces. There is some belief that drinking chilled beverages may be more advantageous, from the standpoint of erosive potential, than drinking warm beverages; as the reduced temperature may favorably alter the kinetics associated with erosive acid attack and the resulting insult to the enamel surface.³⁸ It is probably best to advise patients to wait 1-2 hours after ingesting acidic products before brushing; and when they do brush, it is best to use a stabilized SnF₂ dentifrice that has been demonstrated effective in its ability to help protect against erosive acid damage.

From the professional standpoint, there are a few key points to consider (Table 10). Clearly, consumer trends are creating new dental needs. Dental health is a constantly evolving issue, and one that requires an eye toward the future in order to help stop problems before they get to epidemic levels. Based on a host of information available from studies conducted in the UK and Europe, as well as initial studies done in the United States, dental erosion is clearly one of the next big trends that is emerging and will require dental intervention. Due to the irreversible nature of dental erosion, this may ultimately be a bigger issue than caries. Caries formation is a reversible process that does not involve, at least in the early stages, total destruction of the enamel crystal matrix. Of primary importance in controlling dental erosion is the recognition and appreciation for preventive measures to be put in place at the earliest ages, in order to preserve the natural enamel surface as long as possible. Particularly important is the transition stage from deciduous to permanent dentition,

where patients need to be reminded of the long-term need to maintain strong, healthy teeth for life.

Conclusions

While caries and dental erosion involve the loss of mineral, there are differences between caries and erosive processes. Caries occurs under plaque and is the direct result of bacterial acids. The primary acid that causes caries is lactic acid, a byproduct of the breakdown of fermentable carbohydrates (primarily sugar) by plaque bacteria. Erosion, on the other hand, is a result of the direct action of extrinsic, dietary acids; such as those found in carbonated drinks and fruit juices or intrinsic acids, such as from GERD. With caries, the mineral structure remains intact. Thus, fluoride and other mineral are able to penetrate into the enamel crystal matrix and rebuild or remineralize the challenged enamel. However, dental erosion is different. Once erosive factors overwhelm the pellicle, the result is an initial, relatively fast softening of the enamel followed by abrasive insults that result in complete and permanent removal of the enamel crystal. Net, there is no crystal structure to rebuild.

Dental erosion, a major component under the umbrella term of ETW, is multifactorial and its prevalence is increasing, especially in adolescents and older adults. Advanced ETW causes patients to experience problems with esthetics, function, and pain, and creates treatment dilemmas for dental professionals. Effective management of ETW includes screening and evaluation of all etiological factors, preventive and restorative care, and using the least invasive therapy possible.

Dentifrices containing stabilized SnF₂ have been shown to be very effective at inhibiting both the initiation and progression of dental erosion. ETW must be effectively managed, with a focus on preventive care at the earliest

stages, and monitoring and evaluation of ETW management should be performed regularly during recall sessions. This will help reduce the need for extensive and expensive restorative care in the future.

Table 10. Key areas of erosion awareness for dental professionals.







Erosion Awareness for Professionals.	
Consumer trends are creating new dental needs.	
Dental erosion is an emerging trend.	
Maintaining strong, healthy teeth for life begins at the earliest ages.	
Dental erosion is irreversible.	

Table 10. Key areas of erosion awareness for dental professionals.

<p>Fluoride provides some protection from erosion, but all fluorides are not equally effective.</p>	
<p>SnF_2 is recognized as being highly effective in the prevention of dental erosion.</p>	

Course Test Preview

To receive Continuing Education credit for this course, you must complete the online test. Please go to: www.dentalcare.com/en-us/professional-education/ce-courses/ce517/start-test

- 1. Which of the following statements about the erosion versus caries processes is false?**
 - A. Erosion occurs on the surface; caries development begins sub-surface.
 - B. Caries occurs on the surface, while erosion is a subsurface phenomenon.
 - C. Erosion is often the result of excessive ingestion of acidic beverages.
 - D. Unlike caries, which is a result of bacterial acids, erosion is a result of external acids.
- 2. Methods that have proven useful for measuring the progression of erosive lesions in the laboratory include _____.**
 - A. microradiography
 - B. contact profilometry
 - C. confocal laser scanning microscopy
 - D. Only A and B
 - E. A, B and C
- 3. Which ingredient has the most evidence behind it demonstrating benefits to prevent erosive acid damage?**
 - A. Sodium fluoride.
 - B. Sodium monofluorophosphate.
 - C. Stannous fluoride.
 - D. All of the above.
 - E. A and B.
- 4. Over time, the difference in erosion prevention effectiveness of SnF₂ over NaF, as measured in human in situ erosion prevention studies, becomes _____.**
 - A. more apparent
 - B. less apparent
- 5. The severity of erosive acid attacks depends on which of the following factors?**
 - A. pH
 - B. Titratable acidity
 - C. Buffer capacity
 - D. Contact time on the teeth
 - E. All of the above.
- 6. Preventive habits that reduce the risk of ETW include _____.**
 - A. staying hydrated
 - B. rinsing with water before brushing
 - C. brushing with a stannous fluoride toothpaste
 - D. not brushing for at least 1-2 hours after an acid challenge
 - E. All of the above.
- 7. Which of the following low pH foods or beverages is not considered to be highly erosive?**
 - A. Acid-containing vegetables
 - B. Fruit juices
 - C. Yogurt
 - D. Fresh fruits

8. **Erosive Tooth Wear (ETW) is an umbrella term that includes _____.**
- A. dental erosion
 - B. abfraction
 - C. attrition
 - D. abrasion
 - E. All of the above.
9. **Gastric acids are never associated with erosive tooth wear (ETW).**
- A. False
 - B. True
 - C. It depends on which gastric acid is being considered.
10. **The increasing prevalence of dental erosion is often related to significant increases in the consumption of _____.**
- A. acid containing beverages
 - B. healthier foods, such as fruits and some vegetables
 - C. bottled water
 - D. A and B
 - E. A, B and C
11. **Erosive tooth wear (ETW) is a multifactorial process that may include acid excesses and _____.**
- A. abrasion alone
 - B. attrition alone
 - C. the functional forces of abrasion and attrition
 - D. None of the above.
12. **Which of the following statements best describe acid erosion?**
- A. Dissolution of tooth surfaces by bacterial acids.
 - B. Dissolution of tooth surfaces by acids that are not of biological origin.
 - C. Dissolution of tooth surfaces by either dietary or gastric acids.
 - D. B and C
13. **BEWE is an acronym for _____.**
- A. Begin Erosion Wear Experiment
 - B. Basic Erosive Wear Exam
 - C. Basic Enamel Wasting Estimate
 - D. Biological Enamel Wear Evaluation
14. **Which of the following dietary acids have erosive potential?**
- A. Citric acid
 - B. Phosphoric acid
 - C. Tartaric acid
 - D. Acetic acid
 - E. All of the above.
15. **Dental erosion occurs when the pellicle, nature's natural protection against erosion, is _____.**
- A. overwhelmed by excessive exposure to dietary or gastric acids
 - B. supersaturated with calcium and phosphate from saliva
 - C. heavily stained
 - D. B and C

- 16. The process of dental erosion can be described by which of the following?**
- A. Surface removal due to abrasion of the sound tooth surface, followed by remineralization.
 - B. Surface softening due to erosive acid attack, followed by abrasive factors that can remove this softened layer, followed by additional softening.
 - C. Surface softening due to bacterial acid attack, followed by abrasive factors that can remove this softened layer, followed by additional softening.
 - D. Sub-surface demineralization of tooth mineral, ultimately leading to cavitation.
- 17. Which of the following statements is/are true?**
- A. Erosive tooth surface loss is a growing problem that is seen day-to-day in general practice.
 - B. Both enamel and dentin are susceptible to erosive tooth surface loss.
 - C. BEWE is a simple, quick index for screening a patient's erosion status.
 - D. A, B and C
 - E. Only A and C
- 18. Which of the following statements is/are false?**
- A. Both enamel and dentin are susceptible to erosive tooth surface loss.
 - B. Recent studies indicate the average person in the United States consumes approximately 20 gallons of sugar sweetened beverages each year.
 - C. Dental erosion is increasing in children, adolescents and adults.
 - D. A and C
- 19. Stannous fluoride is more protective than other fluoride sources against dental erosion because _____.**
- A. it deposits a protective, acid resistant layer on exposed tooth surfaces
 - B. it penetrates deeper into the tooth than other fluoride sources
 - C. it tastes better than other fluoride agents
 - D. it has been in use longer than other fluoride actives
- 20. Important aspects of managing patients with erosive tooth wear are _____.**
- A. be familiar with and recognize tooth wear lesions at the earliest possible stage
 - B. implement preventive and treatment measures to preserve the tooth
 - C. reconstruction
 - D. A and B
 - E. A and C
- 21. The most important biological risk factor in Erosive Tooth Wear (ETW) is _____.**
- A. the number of cavities the patient has
 - B. the type of bacteria present in the patient's mouth
 - C. saliva
 - D. the age of the patient
- 22. Which of the following is a primary causative factor for dental erosion?**
- A. *S. mutans*
 - B. *Lactobacillus*
 - C. Sucrose
 - D. *S. sobrinus*
 - E. None of the above.

- 23. Although the prevalence of dental erosion has been reported to be between 7 - 74%, what is the generally accepted value describing the prevalence of dental erosion in teenagers and young adults?**
- A. 10%
 - B. 20%
 - C. 30%
 - D. 40%
- 24. The BEWE index divides the mouth into how many areas for evaluation?**
- A. 2
 - B. 4
 - C. 6
 - D. 8
- 25. Besides saliva, what is another key biological factor that can influence erosion?**
- A. Level of stain.
 - B. Hardness of the teeth.
 - C. The salivary pellicle.
 - D. The number of permanent teeth with cavities.

References

1. O'Brian M. Children's Dental Health in the United Kingdom. Office of Population Censuses and Surveys. London. Her Majesty's Stationary Office. 1994.
2. Nunn JH, Gordon PH, Morris AJ, et al. Dental erosion - changing prevalence? A review of British National childrens' surveys. *Int J Paediatr Dent*. 2003 Mar;13(2):98-105.
3. Linkosalo E, Markkanen H. Dental erosions in relation to lactovegetarian diet. *Scand J Dent Res*. 1985 Oct;93(5):436-41.
4. Wiktorsson AM, Zimmerman M, Angmar-Månsson B. Erosive tooth wear: prevalence and severity in Swedish winetasters. *Eur J Oral Sci*. 1997 Dec;105(6):544-50.
5. Lussi A, Schaffner M, Hotz P, et al. Dental erosion in a population of Swiss adults. *Community Dent Oral Epidemiol*. 1991 Oct;19(5):286-90.
6. Wiegand A, Muller J, Werner C, et al. Prevalence of erosive tooth wear and associated risk factors in 2-7-year-old German kindergarten children. *Oral Dis*. 2006;12(2):117-124. DOI: 10.1111/j.1601-0825.2005.01167.x.
7. Ganss C, Klimek K, Giese K. Dental erosion in children and adolescents--a cross-sectional and longitudinal investigation using study models. *Community Dent Oral Epidemiol*. 2001 Aug;29(4):264-71.
8. Lussi A, Carvalho TS. Erosive tooth wear: a multifactorial condition of growing concern and increasing knowledge. *Monogr Oral Sci*. 2014;25:1-15. doi: 10.1159/000360380. Epub 2014 Jun 26.
9. Luo Y, Zeng XJ, Du MQ, et al. The prevalence of dental erosion in preschool children in China. *J Dent*. 2005 Feb;33(2):115-21. DOI: 10.1016/j.jdent.2004.08.007.
10. Vieira A, Overweg E, Ruben JL, et al. Toothbrush abrasion, simulated tongue friction and attrition of eroded bovine enamel in vitro. *J Dent*. 2006 May;34(5):336-42. Epub 2005 Sep 19. DOI: 10.1016/j.jdent.2005.07.010.
11. Wetton S, Hughes J, West N, et al. Exposure time of enamel and dentine to saliva for protection against erosion: a study in vitro. *Caries Res*. 2006;40(3):213-7. DOI: 10.1159/000092228.
12. Deery C, Wagner ML, Longbottom C, et al. The prevalence of dental erosion in a United States and a United Kingdom sample of adolescents. *Pediatr Dent*. 2000 Nov-Dec;22(6):505-10.
13. Centerwall BC, Armstrong CW, Funkhouser LS, et al. Erosion of dental enamel among competitive swimmers at a gas-chlorinated swimming pool. *Am J Epidemiol*. 1986 Apr;123(4):641-7.
14. Addy M, Shellis RP. Interaction between attrition, abrasion and erosion in tooth wear. *Monogr Oral Sci*. 2006;20:17-31. DOI: 10.1159/000093348.
15. Jaeggi T, Lussi A. Prevalence, incidence and distribution of erosion. *Monogr Oral Sci*. 2014;25:55-73. doi: 10.1159/000360973. Epub 2014 Jun 26. DOI: 10.1159/000360973.
16. Wu B, Liang J, Plassman BL, et al. Edentulism trends among middle-aged and older adults in the United States: Comparison of five racial/ethnic groups. *Community Dent Oral Epidemiol*. 2012 Apr;40(2):145-53. doi: 10.1111/j.1600-0528.2011.00640.x. Epub 2011 Oct 6.
17. Ganss C. Definition of erosion and links to tooth wear. *Monogr Oral Sci*. 2006;20:9-16. DOI: 10.1159/000093344.
18. Lussi A, Schlueter N, Rakhmatullina E, et al. Dental Erosion – An Overview with Emphasis on Chemical and Histopathological Aspects. *Caries Res*. 2011;45 Suppl 1:2-12. doi: 10.1159/000325915. Epub 2011 May 31.
19. Buzalaf MA, Kato MT, Hannas AR. The role of matrix metalloproteinases in dental erosion. *Adv Dent Res*. 2012 Sep;24(2):72-6. doi: 10.1177/0022034512455029.
20. Lussi A, Schaffner M. Progression of and risk factors for dental erosion and wedge-shaped defects over a 6-year period. *Caries Res*. 2000 Mar-Apr;34(2):182-7. DOI: 16587.
21. Hughes JA, West NX, Parker DM, et al. Development and evaluation of a low erosive blackcurrant juice drink in vitro and in situ 1. Comparison with orange juice. *J Dent*. 1999 May;27(4):285-9.

22. Hughes JA, West NX, Parker DM, et al. Development and evaluation of a low erosive blackcurrant juice drink. 3. Final drink and concentrate, formulae comparisons in situ and overview of the concept. *J Dent.* 1999 Jul;27(5):345-50.
23. Calvadini C, Siega-Riz AM, Popkin BM. US adolescent food intake trends from 1965 to 1996. *Arch Dis Child.* 2000 Jul;83(1):18-24.
24. Gleason P, Sutor C. Children's diets in the mid-1990s: Dietary intake and its relationship with school meal participation. Princeton, NJ. Mathematica Policy Research, Inc. 2001.
25. Credit Suisse. Research Institute. Sugar Consumption at a crossroads. September 2013. Accessed March 28, 2017.
26. Per capita consumption of soft drinks in the United States from 2010 to 2014 (in gallons). Statista. Accessed March 28, 2017.
27. Magalhães AC, Moraes SM, Rios D, et al. Effect of ion supplementation of a commercial soft drink on tooth enamel erosion. *Food Addit Contam Part A Chem Anal Control Expo Risk Assess.* 2009 Feb;26(2):152-6. doi: 10.1080/02652030802425326.
28. Richards D. Impact of diet on tooth erosion. *Evid Based Dent.* 2016 Jun;17(2):40. doi: 10.1038/sj.ebd.6401164.
29. Zero DT, Lussi A. Behavioral Factors. *Monogr Oral Sci.* 2006;20:100-05. doi: 10.1159/000093356.
30. Jain P, Nihill P, Sobkowski J, et al. Commercial soft drinks: pH and in vitro dissolution of enamel. *Gen Dent.* 2007 Mar-Apr; 55(2):150-154.
31. Kanzow P, Wegehaupt FJ, Attin T, et al. Etiology and pathogenesis of dental erosion. *Quintessence Int.* 2016 Apr;47(4):275-8. doi: 10.3290/j.qi.a35625.
32. Salas MM, Nascimento GG, Vargas-Ferreira F, et al. Diet influenced tooth erosion prevalence in children and adolescents: Results of a meta-analysis and meta-regression. *J Dent.* 2015 Aug;43(8):865-75. doi: 10.1016/j.jdent.2015.05.012. Epub 2015 Jun 7.
33. Lussi A, Jaeggi T, Zero D. The role of diet in the aetiology of dental erosion. *Caries Res.* 2004;38 Suppl 1:34-44. DOI: 10.1159/000074360.
34. Aykut-Yetkiner A, Wiegand A, Ronay V, et al. In vitro evaluation of the erosive potential of viscosity-modified soft acidic drinks on enamel. *Clin Oral Investig.* 2014 Apr;18(3):769-73. doi: 10.1007/s00784-013-1037-9. Epub 2013 Jul 28.
35. Aykut-Yetkiner A, Wiegand A, Bollhalder A, et al. Effect of acidic solution viscosity on enamel erosion. *J Dent Res.* 2013 Mar;92(3):289-94. doi: 10.1177/0022034512473115. Epub 2013 Jan 11.
36. Hara AT, Lussi A, Zero DT. Biological factors. *Monogr Oral Sci.* 2006;20:88-99. DOI: 10.1159/000093355.
37. Hellwig E, Lussi A. Oral hygiene products and acidic medicines. *Monogr Oral Sci.* 2006;20:112-8. DOI: 10.1159/000093358.
38. Shellis RP, Featherstone JD, Lussi A. Understanding the chemistry of dental erosion. *Monogr Oral Sci.* 2014;25:163-79. Doi: 10.1159/000359943. Epub 2014 Jun 26.
39. Hannig M, Fiebiger M, Güntzer M, et al. Protective effect of the in situ formed short-term salivary pellicle. *Arch Oral Biol.* 2004 Nov;49(11):903-10. DOI: 10.1016/j.archoralbio.2004.05.008.
40. Moazzez R, Bartlett D. Intrinsic causes of erosion. *Monogr Oral Sci.* 2014;25:180-96. doi: 10.1159/000360369. Epub 2014 Jun 26. DOI: 10.1159/000360369.
41. Shah NH, LePendu P, Bauer-Mehren A, et al. Proton pump inhibitor usage and the risk of myocardial infarction in the general population. *PLoS One.* 2015 Jun 10;10(6):e0124653. doi: 10.1371/journal.pone.0124653. eCollection 2015.
42. Fass R, Dickman R. Clinical consequences of silent gastroesophageal reflux disease. *Curr Gastroenterol Rep.* 2006 Jun;8(3):195-201.
43. Wilder-Smith CH, Wilder-Smith P, Kawakami-Wong H, et al. Quantification of dental erosions in patients with GERD using optical coherence tomography before and after double-blind, randomized treatment with esomeprazole or placebo. *Am J Gastroenterol.* 2009 Nov;104(11):2788-95. doi: 10.1038/ajg.2009.441. Epub 2009 Aug 4.
44. Corica, Caprioglio A. Meta-analysis of the prevalence of tooth wear in primary dentition. *Eur J Paediatr Dent.* 2014 Dec;15(4):385-8.

45. Johnson NW. Risk markers for oral diseases. Vol. 1, Dental caries : markers of high and low risk groups and individuals. Cambridge. Cambridge University Press. 1991. 266-285.
46. Loesche WJ. Role of streptococcus mutans in dental decay. *Microbiol Rev.* 1986 Dec;50(4):353-80.
47. Featherstone JD. Dental caries: a dynamic disease process. *Aust Dent J.* 2008 Sep;53(3):286-91. doi: 10.1111/j.1834-7819.2008.00064.x.
48. Centers for Disease Control and Prevention (CDC). Ten Great Public Health Achievements – United States, 1990-1999. *MMWR Morb Mortal Wkly Rep.* 1999 Apr 2;48(12):241-3.
49. Featherstone JD. Prevention and reversal of dental caries: role of low level fluoride. *Community Dent Oral Epidemiol.* 1999 Feb;27(1):31-40.
50. Ganss C, Lussi A, Schlueter N. The histological features and physical properties of eroded dental hard tissues. *Monogr Oral Sci.* 2014;25:99-107. doi: 10.1159/000359939. Epub 2014 Jun 26.
51. Eversole SL, Saunders-Burkhardt K, Faller RV. Erosion protection comparison of stabilised SnF₂, mixed fluoride active and SMFP/arginine-containing dentifrices. *Int Dent J.* 2014 Mar;64 Suppl 1:22-8. doi: 10.1111/idj.12099.
52. Faller RV, Eversole SL. Enamel protection from acid challenge – benefits of marketed fluoride dentifrices. *J Clin Dent.* 2013;24(1):25-30.
53. Faller RV, Eversole SL, Tzeghai GE. Enamel protection: a comparison of marketed dentifrice performance against dental erosion. *Am J Dent.* 2011 Aug;24(4):205-10.
54. Eversole SL, Saunders-Burkhardt K, Faller RV. Erosion prevention potential of an over-the-counter stabilized SnF₂ dentifrice compared to 5000ppm F prescription strength products. *J Clin Dent.* 2015;26(2):44-9.
55. Hooper SM, Newcombe RG, Faller R, et al. The protective effects of toothpaste against erosion by orange juice: studies in situ and in vitro. *J Dent.* 2007 Jun;35(6):476-81. Epub 2007 Feb 27. DOI: 10.1016/j.jdent.2007.01.003.
56. Huysmans MCDNJM, Jager DHJ, Ruben JL, et al. Reduction of erosive wear in situ by stannous fluoride-containing toothpaste. *Caries Res.* 2011;45(6):518-23. doi: 10.1159/000331391. Epub 2011 Oct 5.
57. Bellamy PG, Harris R, Date RF, et al. In situ clinical evaluation of a stabilised, stannous fluoride dentifrice. *Int Dent J.* 2014 Mar;64 Suppl 1:43-50. doi: 10.1111/idj.12102.
58. West NX, Seong J, Hellin N, et al. A clinical study to measure anti-erosion properties of a stabilized stannous fluoride dentifrice relative to a sodium fluoride/triclosan dentifrice. *Int J Dent Hyg.* 2015 Jun 22. doi: 10.1111/idh.12159. [Epub ahead of print].
59. West NX, He T, Macdonald EL, et al. Erosion protection benefits of stabilized SnF₂ dentifrice versus an arginine – sodium monofluorophosphate dentifrice: Results from in vitro and in situ clinical studies. *Clin Oral Investig.* 2017 Mar;21(2):533-540. doi: 10.1007/s00784-016-1905-1. Epub 2016 Aug 1.
60. West N, Seong J, Macdonald E, et al. A randomised clinical study to measure the anti-erosion benefits of a stannous-containing sodium fluoride dentifrice. *J Indian Soc Periodontol.* 2015 Mar-Apr;19(2):182-7. doi: 10.4103/0972-124X.145817.
61. Khambe D, Eversole SL, Mills T, et al. Protective effects of SnF₂ – Part II. Deposition and retention on pellicle-coated enamel. *Int Dent J.* 2014 Mar;64 Suppl 1:11-5. doi: 10.1111/idj.12097.
62. Faller RV. Meeting the challenges of tooth sensitivity and dental erosion with stannous fluoride. *Cosmetics & Toiletries.* 2012; 127(5):362-371.
63. Paepegaey AM, Barker ML, Bartlett DW, et al. Measuring enamel erosion: a comparative study of contact profilometry, non-contact profilometry and confocal laser scanning microscopy. *Dent Mater.* 2013 Dec;29(12):1265-72. doi: 10.1016/j.dental.2013.09.015. Epub 2013 Oct 24.
64. Bartlett DW, Lussi A, West NX, et al. Prevalence of tooth wear on buccal and lingual surfaces and possible risk factors in young European adults. *J Dent.* 2013 Nov;41(11):1007-13. doi: 10.1016/j.jdent.2013.08.018. Epub 2013 Sep 1.
65. Carvalho TS, Lussi A, Jaeggi T, et al. Erosive tooth wear in children. *Monogr Oral Sci.* 2014;25:262-78. doi: 10.1159/000360712. Epub 2014 Jun 26.

66. Carvalho TS, Colon P, Ganss C, et al. Consensus report of the European Federation of Conservative Dentistry: erosive tooth wear--diagnosis and management. *Clin Oral Investig*. 2015 Sep;19(7):1557-61. doi: 10.1007/s00784-015-1511-7. Epub 2015 Jul 1.
67. Ganesh M, Hertzberg A, Nurko S, et al. Acid rather than nonacid reflux burden is a predictor of tooth erosion. *J Pediatr Gastroenterol Nutr*. 2016 Feb;62(2):309-13. doi: 10.1097/MPG.0000000000000927.
68. Bartlett DW, Ganss C, Lussi A. Basic Erosive Wear Examination (BEWE): a new scoring system for scientific and clinical needs. *Clin Oral Investig*. 2008 Mar;12 Suppl 1:S65-8. doi: 10.1007/s00784-007-0181-5. Epub 2008 Jan 29.
69. Van der Weijden FA, Campbell SL, Dörfer CE, et al. Safety of oscillating-rotating powered brushes compared to manual toothbrushes: a systematic review. *J Periodontol*. 2011 Jan;82(1):5-24. doi: 10.1902/jop.2010.100393. Epub 2010 Sep 10.
70. Wiegand A, Burkhard JP, Eggmann F, et al. Brushing force of manual and sonic toothbrushes affects dental hard tissue abrasion. *Clin Oral Investig*. 2013 Apr;17(3):815-22. doi: 10.1007/s00784-012-0788-z. Epub 2012 Jul 13.
71. Hara AT, Ando M, Gonzales-Cabezas C, et al. Protective effect of the dental pellicle against erosive challenges in situ. *J Dent Res*. 2006 Jul;85(7):612-6.

About the Authors

Donald J. White, PhD



Dr. Donald White is a Research Fellow at the Procter & Gamble Company Health Care Research Center in Cincinnati, Ohio, USA. Dr. White received his BS in Chemistry and PhD in Physical Chemistry from the State University of New York at Buffalo, carrying out research on the mechanisms of formation and solubilization of kidney stones. Since 1982, Dr. White's dental research interests include the contribution of physical chemistry to Oral Biological processes including fundamental aspects of tartar formation, stain development, dental caries, and dental plaque formation. Dr. White has published extensively, including over 100 peer-reviewed publications, approximately 200 meeting presentations and abstracts, and over 40 global patents.

Email: white.dj.1@pg.com

Warden H. Noble, DDS, MS, MEd



Dr. Noble graduated from and was on the faculty at University of California, San Francisco. He is a board certified prosthodontist and was formerly a Clinical Professor and Co-director of Graduate Prosthodontics at UCSF. After 40 years in private practice, he is now a Professor in the Department of Integrated Reconstructive Dental Sciences at the Arthur A. Dugoni School of Dentistry, University of the Pacific, San Francisco, CA.

Email: wnoble@pacific.edu

Nicola West, BDS, FDSRCS, PhD FDS (Rest Dent)



Professor Nicola West is a graduate of the University of Wales College of Medicine (Cardiff). After attaining her fellowship from the Royal College of Surgeons of England, Nicola gained her PhD from the University of Wales (Cardiff) in 1995, on dentine hypersensitivity and tooth surface loss. Nicola gained her chair in Periodontology at Bristol Dental School, where she conducts research at an international level, leading the Clinical Trials Unit and attracting substantial industrial funding. Current research interests include: developing research methodologies, periodontal conditions, bone augmentation and peri-implant lesions, the scientific evaluation of oral health care products, tooth wear, dentine hypersensitivity and tooth whitening. Nicola leads Restorative Dentistry at the Bristol Dental School and is an Honorary Consultant.

Nicola lectures regularly in the national and international forum, is an active member and secretary of the British Society of Periodontology and currently is the regional postgraduate tutor for the British Society of Periodontology in the South West of England and European BSP representative. She is a Fellow of the Higher Education Academy, a UK member of the International Standards Committee and on the advisory board of the British Dental Health Foundation.

Email: N.X.West@bristol.ac.uk